



## OPEN ACCESS

## EDITED BY

Alex Rafacho,  
Federal University of Santa Catarina,  
Brazil

## REVIEWED BY

Mustafa Oztop,  
Mehmet Akif Ersoy University, Türkiye  
Shannon Noella Tessier,  
Massachusetts General Hospital and  
Harvard Medical School, United States

## \*CORRESPONDENCE

Scott T. Cooper,  
✉ scooper@uwf.edu

RECEIVED 16 April 2023

ACCEPTED 12 June 2023

PUBLISHED 26 June 2023

## CITATION

De Vrij EL, Bouma HR, Henning RH and  
Cooper ST (2023), Hibernation  
and hemostasis.  
*Front. Physiol.* 14:1207003.  
doi: 10.3389/fphys.2023.1207003

## COPYRIGHT

© 2023 De Vrij, Bouma, Henning and  
Cooper. This is an open-access article  
distributed under the terms of the  
[Creative Commons Attribution License  
\(CC BY\)](#). The use, distribution or  
reproduction in other forums is  
permitted, provided the original author(s)  
and the copyright owner(s) are credited  
and that the original publication in this  
journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Hibernation and hemostasis

Edwin L. De Vrij<sup>1,2</sup>, Hjalmar R. Bouma<sup>2,3</sup>, Robert H. Henning<sup>2</sup> and  
Scott T. Cooper<sup>4\*</sup>

<sup>1</sup>Department of Plastic Surgery, University Medical Center Groningen, University of Groningen, Groningen, Netherlands, <sup>2</sup>Department of Clinical Pharmacy and Pharmacology, University Medical Center Groningen, Groningen, Netherlands, <sup>3</sup>Department of Internal Medicine, University Medical Center Groningen, University of Groningen, Groningen, Netherlands, <sup>4</sup>Biology Department, University of Wisconsin-La Crosse, La Crosse, WI, United States

Hibernating mammals have developed many physiological adaptations to accommodate their decreased metabolism, body temperature, heart rate and prolonged immobility without suffering organ injury. During hibernation, the animals must suppress blood clotting to survive prolonged periods of immobility and decreased blood flow that could otherwise lead to the formation of potentially lethal clots. Conversely, upon arousal hibernators must be able to quickly restore normal clotting activity to avoid bleeding. Studies in multiple species of hibernating mammals have shown reversible decreases in circulating platelets, cells involved in hemostasis, as well as in protein coagulation factors during torpor. Hibernator platelets themselves also have adaptations that allow them to survive in the cold, while those from non-hibernating mammals undergo lesions during cold exposure that lead to their rapid clearance from circulation when re-transfused. While platelets lack a nucleus with DNA, they contain RNA and other organelles including mitochondria, in which metabolic adaptations may play a role in hibernator's platelet resistance to cold induced lesions. Finally, the breakdown of clots, fibrinolysis, is accelerated during torpor. Collectively, these reversible physiological and metabolic adaptations allow hibernating mammals to survive low blood flow, low body temperature, and immobility without the formation of clots during torpor, yet have normal hemostasis when not hibernating. In this review we summarize blood clotting changes and the underlying mechanisms in multiple species of hibernating mammals. We also discuss possible medical applications to improve cold preservation of platelets and antithrombotic therapy.

## KEYWORDS

hibernation, torpor, hemostasis, platelet, coagulation, metabolism, hypothermia

## Overview of hemostasis

Blood plays an essential role in vertebrates by circulating oxygen and nutrients and removing wastes from all tissues. This requires an extensive network of vessels that contain fluid blood, yet can form solid clots if vessels are damaged to prevent the organism from bleeding to death. Spontaneous blot clot formation is counterbalanced by anticoagulant mechanisms that can become overwhelmed at the site of a wound leading to hemostasis. The word hemostasis is derived from the Greek αίμα/hema (=blood) and στάσις/stasis (=halt), literally the stopping of blood. Hemostasis is generally divided into three phases, primary hemostasis involving anucleated cells called platelets, secondary hemostasis triggered by a cascade of serine proteases to form a fibrin clot, and fibrinolysis, which breaks down the clot (Figure 1). There are multiple ways to activate hemostasis including endothelial cell injury, clotting factors or platelet alterations, and abnormal blood flow. During torpor, heart rate


and blood flow are reduced to levels that would lead to the formation of potentially lethal blood clots in most non-hibernating mammals. Hypothermia also affects both procoagulant and anticoagulant aspects of hemostasis. Hibernating mammals thus need multiple adaptations to reversibly suppress blood clotting during torpor, which will be discussed in this review.


## Hibernating mammalian models

Hibernators encounter several risk factors for thrombosis including low body temperature, stasis of blood, hypercoagulability, and endothelial activation. During torpor a ground squirrel's body temperature drops from 35°C–38°C to 4°C–8°C (Lechler and Penick, 1963; Reddick et al., 1973), the heart rate is reduced from its normal 200–300 to three to five beats/min (Zatzman, 1984), and respiration is reduced from 100 to 200 to four to six breaths/min (McArthur and Milsom, 1991). As expected, blood pressure also drops from 140/100 mmHg to 60/30 mmHg, with values as low as 10 mmHg reported (Lyman and O'Brien 1960). Thus hibernation entails periods of prolonged immobility (Carey et al., 2003; Utz et al., 2009) with low blood flow (stasis) in veins and atria (Horwitz et al., 2013),

increased blood viscosity (Kirkebo, 1968; Halikas and Bowers, 1973; Arinell et al., 2018), cycles of hypoxia-reoxygenation, and cooling-rewarming with signs of endothelial activation (Carey et al., 2003; Talaei et al., 2012). Additionally, at entrance of the hibernation season, hibernators are generally grossly overweight (Martin, 2008). Remarkably, despite this multitude of risk factors for thromboembolism, hibernators do not demonstrate signs of organ damage due to thromboembolic complications during hibernation or upon arousal in spring (Cooper et al., 2012; Bonis et al., 2019).

A reversible decrease in hemostasis has been demonstrated during torpor in diverse hibernating animals. In small mammalian deep hibernators body temperature can drop to 2°C–4°C as found in ground squirrels (Svihla et al., 1951; Lechler and Penick, 1963; Reddick et al., 1973; Pivrun and Sinnamon, 1981; Bouma et al., 2010; Cooper et al., 2012; Hu et al., 2017; Öztöp et al., 2019), hamsters (Denyes and Carter, 1961; Deveci et al., 2001; de Vrij et al., 2014), hedgehogs (Biorck et al., 1962; De Wit et al., 1985), and bats (Rashid et al., 2016). In contrast, large mammals like bears show more moderate body temperatures during hibernation by only dropping from 37°C to 32°C coupled with prolonged immobility (Fröbert et al., 2010; Welinder et al., 2016; Iles et al., 2017; Arinell et al., 2018; Thienel et al., 2023). Decreased hemostasis in





hibernators may partly be explained by the decrease in body temperature, as forced hypothermia also lowers platelet counts in hamsters, and even suppresses hemostasis in non-hibernating species like rats and mice (de Vrij et al., 2014). Finally, ectothermic vertebrates also reduce hemostasis during hibernation as seen in turtles (Barone and Jacques, 1975) and frogs (Ahmad et al., 1979). Given that a wide range of hibernating species suppress their hemostasis during torpor through multiple mechanisms, it is likely that this represents a vital physiological adaptation to multiple risk factors during torpor. The rapid restoration of hemostasis after arousal indicates that being able to form clots soon after arousal is also imperative. Changes in clot formation throughout hibernation is illustrated graphically by thromboelastograms where whole blood is treated with an activator of coagulation and the speed and strength of clot formation is measured (Figure 2). Because both primary and secondary hemostasis are regulated by proteins in circulation and platelets which lack a nucleus, the adaptations to rapid changes in temperature and blood flow must be inherent in these proteins and cells, and not rely on new synthesis as seen in other organs.


## Effect of hypothermia on hemostasis

While clot formation may prevent bleeding after tissue damage and initiate wound healing, inadvertent generation of a thrombus can be detrimental for any organism. Virchow's Triad was first described in 1856 and identified venous stasis, vascular injury, and hypercoagulability as risk factors for thrombosis. The ensuing

disseminated intravascular coagulation (DIC) of hypothermic patients may result in consumption of clotting factors, ischemia, and necrosis of organs and eventually result in death (Duguid et al., 1961; Stine, 1977; Mahajan et al., 1981). Besides provoking prothrombotic effects, decreased body temperature slows the enzymatic reactions in secondary hemostasis (Rohrer and Natale, 1992; DeLong et al., 2017), prolonging bleeding time in cold skin, and diminishing thromboxane A release from platelets (Valeri et al., 1995; Michelson et al., 1999). Consequently, hypothermia is associated with a secondary hypocoagulation state with prolonged time to clot formation in tests such as PT and APTT when temperatures drop below 35°C (Rohrer and Natale, 1992; Polderman, 2009). Low temperature *in vivo* increases activation, aggregation, and reversible storage of platelets in the liver or spleen (Van Poucke et al., 2014). Low temperatures of the extremities have been implicated to 'prime' platelets for activation at these sites most susceptible to bleeding throughout evolutionary history, which also leads to increased clearance of these platelets from circulation (Hoffmeister et al., 2003). Furthermore, both accidental and therapeutic hypothermia are associated with a reduction in platelet count known as thrombocytopenia (Vella et al., 1988; Chan and Beard, 1993; Mikhailidis and Barradas, 1993; Mallet, 2002; Morrell et al., 2008; Jacobs et al., 2013; Wang et al., 2015). Whether this thrombocytopenia in patients can be reversed quickly by rewarming—or whether thrombocytopenia recovers due to *de novo* production and release of platelets from the bone marrow—is still not clear.

*Ex vivo* cooling of platelets from non-hibernating mammals induces platelet shape changes similar to activation of platelets (White and Krivit, 1967; Winokur and Hartwig, 1995; Peerschke et al., 2008; Egidi et al., 2010), while low temperature also increases degranulation of activated platelets and plasma levels of activation products of platelets (Hoffmeister, 2011; Van Poucke et al., 2014). Cooled platelets demonstrate an increased tendency to aggregate (Xavier et al., 2007) and are rapidly cleared from circulation after transfusion (Becker et al., 1973; Berger et al., 1998; Van Poucke et al., 2014). Currently, human platelets are stored at 22°C–24°C room temperature before transfusion which increases risk of bacterial contamination and thus limits shelf-life to only 5–7 days, compared to 40 days for cold stored red blood cells (D'Alessandro et al., 2010). However, even if low body temperature leads to a 10-fold decrease in enzymatic activity and slowing down of clot formation, the prolonged time in torpor would be sufficient for clots to form during torpor. For example, to produce serum in the lab, blood is collected in the absence of an anticoagulant and stored in a refrigerator where it will still form a fibrin clot at 4°C if given enough time. Further unraveling the temperature effects on hemostasis may yield improved understanding of its role in hypothermic patients and in hibernating mammals and potentially disclose new pathways for drug development focused on novel antithrombotic strategies.

Primary hemostasis involves adherence of circulating platelets to damaged endothelium or subendothelial collagen and aggregation between platelets (Figure 1A). Platelets, which bud off from large multinuclear bone marrow cells called megakaryocytes, are 2–5 μm anucleated blood cells that play a major role in hemostasis,



inflammation, bacterial defense and wound regeneration and are even involved in cancer metastasis (Semple et al., 2011; Ware et al., 2013; Leblanc and Peyruchaud, 2016; Fernandez-Moure et al., 2017). Platelets are activated by many molecules including Von Willebrand Factor (VWF), collagen, thrombin, adenosine diphosphate (ADP) and adrenaline (Nieswandt et al., 2011; Versteeg et al., 2013). Activated platelets express membrane proteins including glycoprotein Ib-IX-V (GPIb-IX-V) and P-selectin that bind to activated endothelium or subendothelial collagen through VWF (Nieswandt et al., 2011) (Figure 3). Platelets degranulate upon activation, releasing molecules that stimulate further platelet activation, the coagulation cascade, inflammation, tissue regeneration and bacterial killing (Semple et al., 2011; Fernandez-Moure et al., 2017). More platelets are then recruited to the site of injury (Figure 1B). Platelets adhere to the subendothelial extracellular matrix (ECM) and form a hemostatic plug by aggregating with each other.

## Primary hemostasis adaptations during torpor

Circulating platelet levels decrease about 10-fold in multiple hibernating species during torpor, with a return to normal levels as the animals rewarm during arousals (Figure 4). These dynamics could decrease the risk of a large intravascular clot forming during torpor, while restoring normal clotting activity upon arousal. The

thrombocytopenia is temperature dependent and can be induced in hamsters, rats, and mice with forced chilling (de Vrij et al., 2014). Cold acclimation of rats and hamsters also reduced circulating platelet levels even though the animal's body temperatures did not decrease (Deveci et al., 2001). The rapid return of platelets in circulation after arousal, in the absence of large numbers of circulating immature platelets, suggests that hibernator platelets are stored during the cold and released into circulation during rewarming. Removal of the spleens of Syrian hamsters and 13-lined ground squirrels before or during torpor did not affect the platelet storage and release indicating that spleen is not required for storage as originally proposed (Reddick et al., 1973; Cooper et al., 2012; Cooper et al., 2016; de Vrij et al., 2021). Rather, platelet storage during torpor occurs in the sinusoids of the liver, consistent with this organ being responsible for storing platelets (Cooper et al., 2017; de Vrij et al., 2021). The mechanism of hepatic storage of platelets during torpor is currently unknown, potentially platelet margination may underlie the reversible storage and release in liver (de Vrij et al., 2021). Although platelets cluster together in liver sinusoids during torpor, they do not form irreversible thrombi. Several adaptations may prevent platelet activation during their slumber in cold liver sinusoids. Platelet GpIba receptors can bind to VWF, but the plasma concentration of VWF decreases during torpor in bears, ground squirrels and hamsters, with a selective loss of the thrombogenic high molecular weight multimers (Cooper et al., 2016; Friedrich et al., 2017). Additionally, platelets from torpid ground squirrels bind less VWF and collagen, consistent with decreased GpIba




FIGURE 4

Temperature association with platelet count reduction is consistent for all hibernating mammals studied so far. Platelet count in torpor as percentage of euthermia platelet count was calculated from values of studies summarized in Table 1. If a study did not report euthermic platelet count, literature data were used. Fitted polynomial quadratic curve (black line) with 95% confidence interval (dotted gray line). Pearson's  $r = 0.96$ ,  $p < 0.05$  (Lechler and Penick, 1963; Reddick et al., 1973; Pivorun and Sinnamon, 1981; Bouma et al., 2010; Fröbert et al., 2010; Cooper et al., 2012; de Vrij et al., 2014; Welinder et al., 2016; Cooper et al., 2017; Hu et al., 2017; Arinell et al., 2018; de Vrij et al., 2021).

activity on their surface. Upon arousal, VWF levels return to normal rapidly, possibly by release of stores in platelets or endothelial cells (Cooper et al., 2016). The combination of decreased platelet and VWF levels along with decreased binding of platelets to VWF would all help to suppress activation of primary hemostasis during torpor-induced stasis.

Shape change is another prominent feature of cooled and hibernating platelets. Under normal conditions, platelets have a circumferential microtubule ring that keeps them in a disc conformation. In non-hibernating mammals, chilling platelets leads to depolymerization of their microtubules and the platelets become spherical and do not repolymerize upon warming (White and Krivit, 1967; Winokur and Hartwig, 1995). In contrast, ground squirrels and Syrian hamster platelet microtubules form rods when chilled and can reconstitute the circumferential ring upon rewarming (Cooper et al., 2017; de Vrij et al., 2021). This allows ground squirrel platelets to retain their overall shape after chilling. It is unclear if this rod shape has any physiological or adaptive significance, but it may protect platelets from damage caused by cold storage.

Cold storage of human and mouse platelets damages platelets in several ways, collectively termed cold storage lesions. Sialic acid residues on surface glycoproteins are cleaved leading to the rapid clearance of chilled platelets from circulation by the liver (Hoffmeister et al., 2003; Hoffmeister, 2011). Cold storage also induces intrinsic mitochondrial activated apoptosis in human and mouse platelets (Van Der Wal et al., 2010; Lebois and Josefsson, 2016; Stolla et al., 2020). In contrast, ground squirrel platelets stored *in vitro* in the cold are not cleared rapidly upon re-transfusion (Cooper et al., 2012). Ground squirrel platelets stored in the cold show decreased desialylation of surface glycoproteins and decreased phagocytosis by hepatocytes. They also appear to be resistant to

induction of apoptosis as measured by caspase activation and surface phosphatidylserine on platelet membranes (Splinter et al., 2023). These adaptations of hibernator platelets to cold storage are consistent with their storage and release after weeks of cold storage during torpor and likely add to the rapid restoration of a hibernator's ability to form clots upon arousal.

Platelets are anucleate cells and their protein content is determined primarily by progenitor megakaryocytes in the bone marrow. Surprisingly, a bone marrow transcriptome of samples collected in the fall, winter, and summer from 13-lined ground squirrels revealed no significant differences in any platelet protein transcripts, including P-selectin, membrane receptors, and integrins (Cooper et al., 2016). The platelet proteome collected in the same species and at the same times of the year, also displayed no differences in integrins and signaling proteins, but did reveal significant seasonal differences in other proteins (Cooper et al., 2021). In the summer, platelets have increased heat shock proteins which could help with proper protein folding at warm temperatures. In the fall, as animals are going through short bouts of shallow torpor, pro-inflammatory and clotting proteins become less abundant in platelets. During torpor, more plasma-derived proteins such as albumin and lipoproteins are present in platelets which are known to inhibit apoptosis and thrombosis (Cooper et al., 2021). A very recent platelet proteome study on brown bears showed a 55-fold decrease in the heat shock protein HSP47 (SerpinH1) during hibernation (Thienel et al., 2023). This decrease correlated with a decline in DVT formation during prolonged immobilization, and was also demonstrated in humans, pigs, and knock-out mice. Interestingly, and in sharp contrast, a previous study of 13-lined ground squirrel platelet proteome documented unchanged HSP47 abundance across various phases of hibernation (Cooper et al., 2021). This marked difference in Hsp47 regulation could be due to bears being shallow hibernators and also of greater body mass compared to ground

TABLE 1 Primary hemostasis in hibernation.

Measurement	Euthermia	Torpor	Arousal	Species	References
<b>Whole-blood clotting time</b>					
(sec)	210 ± 76	315 ± 66*		13-lined ground squirrel	Lechler and Penick (1963)
(min)	2.2 ± 0.3	48.0 ± 5.4*	11.5 ± 1.5* <sup>#</sup>	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
(min)	4	11	5	Hedgehog	Biorck et al. (1962)
(sec)	81 ± 12	217 ± 30	164 ± 33	American black bear	Iles et al. (2017)
<b>Thromboelastography</b>					
R (min)	1.6 ± 0.4	16.2 ± 12.3*	4.7 ± 1.2* <sup>#</sup>	13-lined ground squirrel	Cooper et al. (2012)
Alpha (°)	58.8 ± 11.3	6.6 ± 6.2*	17.1 ± 7.4* <sup>#</sup>	13-lined ground squirrel	Cooper et al. (2012)
Maximum amplitude (mm)	47.2 ± 5.2	6.0 ± 6.8*	17.2 ± 10.4* <sup>#</sup>	13-lined ground squirrel	Cooper et al. (2012)
G (dynes/cm <sup>2</sup> )	4.6 ± 1.0	0.3 ± 0.4*	1.1 ± 0.8* <sup>#</sup>	13-lined ground squirrel	Cooper et al. (2012)
<b>Platelet aggregation (arbitrary units)</b>					
ADP	70.0 ± 26.6	29.2 ± 8*		Scandinavian brown bear	Arinell et al. (2018)
	66 ± 23	33 ± 10*		Scandinavian brown bear	Arinell et al. (2018)
Arachidonic acid	73 ± 16	28 ± 9*		Scandinavian brown bear	Arinell et al. (2018)
	68 ± 20	33 ± 10*		Scandinavian brown bear	Arinell et al. (2018)
Collagen	68.3 ± 17	30.7 ± 10		Scandinavian brown bear	Arinell et al. (2018)
	63 ± 22	30 ± 7*		Scandinavian brown bear	Arinell et al. (2018)
TRAP	18.5 ± 10.0	9.2 ± 6.9*		Scandinavian brown bear	Arinell et al. (2018)
PAR-4	22.5 ± 7.1	12.7 ± 7.1*		Scandinavian brown bear	Arinell et al. (2018)
<b>Platelet count</b>					
(x 10 <sup>9</sup> /L)	445.2 ± 123.4 36.3°C	47.9 ± 22.3* 7.9°C		13-lined ground squirrel	Lechler and Penick (1963)
	303.6 ± 10.6 37°C	45 ± 3.4* 6°C	232.4 ± 19.7* <sup>#</sup> 37°C	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
	375.3 ± 40.8 37°C	114.2 ± 36.0* 9°C	217.0 ± 35.9* <sup>#</sup> 37°C	Daurian ground squirrel	Hu et al. (2017)
	293 ± 81 36°C	44 ± 30.9* 8°C	194 ± 5* <sup>#</sup> 35°C	European ground squirrel	Bouma et al. (2010)
		23.3 ± 1.3* 9.8°C	410.9 ± 59.2 36°C	13-lined ground squirrel	Cooper et al. (2012)
	394 ± 157	55 ± 30*		13-lined ground squirrel	Reddick et al. (1973)
	451 ± 87 35.5°C	128 ± 81 9.8°C	498 ± 144 36.4°C	13-lined ground squirrel	Cooper et al. (2017)
	797 ± 124 35°C	381 ± 239* 25°C	739 ± 253 <sup>#</sup> 35°C	Djungarian hamster	de Vrij et al. (2014)
	198 ± 6 35°C	8 * 8°C	187 <sup>#</sup> 35°C	Syrian hamster	de Vrij et al. (2014)
	430 ± 82 35.7°C	36 ± 17* 8.2°C	468 ± 100 35.4°C	Syrian hamster	de Vrij et al. (2021)
	207 ± 24			Scandinavian brown bear	Fröbert et al. (2010)
	262 ± 61 40°C	174 ± 51* 33°C	262 ± 61	Scandinavian brown bear	Arinell et al. (2018)
	229 ± 39	146 ± 47*		Scandinavian brown bear	Arinell et al. (2018)
	228 ± 36 37°C	149 ± 43* 32°C		Scandinavian brown bear	Welinder et al. (2016)

(Continued on following page)

TABLE 1 (Continued) Primary hemostasis in hibernation.


Measurement	Euthermia	Torpor	Arousal	Species	References
(%EU)	100% ± 31% (n = 19) 34°C	65% ± 29%* (n = 12) 26°C	111% ± 26% (n = 13) 33°C	Daily torpor C57Bl/6 Mouse	(De Vrij et al. unpublished)
(10 <sup>3</sup> /mm <sup>3</sup> Mean ± SE)	5.6 ± 0.6		6.0 ± 1.1	Common yellow bat	Rashid et al. (2016)
(10 <sup>3</sup> /mm <sup>3</sup> Mean ± SE)	7.5 ± 1.0		7.4 ± 1.2	Common pipistrelle bat	(Rashid et al., 2016)
<b>P-selectin expressing platelets (%)</b>	8 ± 7	0	7 ± 6	Syrian hamster	de Vrij et al. (2014)
	3 ± 1	11 ± 7	10 ± 3	Syrian hamster	de Vrij et al. (2021)
<b>Platelets activated by ADP (%)</b>	16 ± 14	16 ± 6	29 ± 6	Syrian hamster	de Vrij et al. (2014)
	22 ± 8	16 ± 2	47 ± 11* <sup>a</sup>	Syrian hamster	de Vrij et al. (2021)
<b>VWF (% relative to human plasma)</b>	24.9 ± 3.7	2.4 ± 0.01*		13-lined ground squirrel	Cooper et al. (2016)
(%EU)	100% ± 10.5% (n = 10)	7.3% ± 3.2%* (n = 4)	8.4% ± 11.5%* (n = 4)	Syrian hamster	(De Vrij et al. unpublished)
(IU/mL)	1.7 ± 0.2	1.3 ± 0.2*		Scandinavian brown bear	Welinder et al. (2016)
<b>VWF:collagen binding activity (%EU)</b>	100% ± 10% (n = 7)	35% ± 15% (n = 4)	25% ± 33%* (n = 4)	Syrian hamster	(De Vrij et al. unpublished)

squirrel, so the immobilization puts bears at greater risk of DVT. Moreover, torpid bears have a modest drop in circulating platelets compared to deep hibernators (Figure 4), possibly warranting additional protection against DVT. It would be interesting to look more closely at the inhibition of HSP47 in deep hibernators other than 13 lined ground squirrel to appreciate its role in hibernation. In addition to the decrease in circulating platelet numbers, these proteomic changes could prevent unwanted activation of platelets during stasis or during their storage in liver. While there were no significant differences in the amounts of surface receptors or proteins in signaling pathways at the proteome level, it is possible that the activities of these pathways are altered. Alternatively, some of the proteins may be sequestered in the cytoplasm or secretory granules which could be resolved by immunohistochemical analysis. Studying the phosphoproteome and metabolome of platelets in torpor would help to resolve which pathways are affected.

In recent years, cultured cells from hibernators have demonstrated to hold specific adaptations that confer protection from cooling in a cell-autonomous way (Talaie et al., 2011; Ou et al., 2018; Hendriks et al., 2020). Mechanistically, these studies in hamster cell lines and ground squirrel pluripotent stem cells derived neurons indicate that adaptations in mitochondrial function, allowing cooled cells to maintain ATP production and lower reactive oxygen species production, constitute a key element of cell protection. Given that platelets contain mitochondria, we re-analyzed aforementioned ground squirrel platelet proteome data (Cooper et al., 2021) by separating out mitochondrial proteins. Analysis of the platelet proteins involved in metabolism revealed an increase in 79 proteins with clusters in lipoprotein metabolism, cholesterol esterification, and lipase activity during the winter. With respect to mitochondrial proteins, 24 were increased during the winter relative to summer and fall, with clusters in fatty acid beta-oxidation, the electron transport chain, and protein targeting to the mitochondria (Figure 5). These proteome changes correspond well with changes in most other tissues of hibernators, principally denoting the shift to lipid metabolism during hibernation as animals rely on fat stores for energy. As

platelets appear to be no exception, it is conceivable that they hold similar cell autonomous protection from cooling damage as found in hamster and ground squirrel smooth muscle cells, kidney cells and neurons. Interestingly, ground squirrel neurons maintain their tubulin network during cooling by limitation of the production of reactive oxygen species (Ou et al., 2018). Such mechanism may also support tubulin stability and possibly underlie the difference in cooling induced platelet shape change between hibernators (reversible rod formation) and non-hibernators (irreversible sphere formation). Given this clearcut difference in response to cooling and the ease of obtaining primary cells, platelets constitute an important platform for the future study of cell-autonomous adaptations in hibernators.

Secondary hemostasis occurs simultaneously with primary hemostasis forming a fibrin network to trap red and white blood cells and further strengthen the hemostatic platelet plug (Figure 6). Secondary hemostasis occurs via a plasma coagulation cascade divided into the intrinsic and extrinsic pathways. Most clotting factors are serine proteases that are always present in plasma as inactive zymogens (identified by Roman numerals II, VII, IX, X, XI, and XII) and are activated by proteolysis (addition of an “a”). Factors V and VIII are non-enzymatic cofactors also activated by proteolysis. The extrinsic pathway starts with exposure and binding of tissue factor (TF) to plasma coagulation factor VII, which forms a TF/VIIa complex. The TF/VIIa complex proteolytically activates factors IX and X, commencing the common pathway, creating a prothrombinase complex with Va that converts prothrombin (factor II) into thrombin (IIa) (Versteeg et al., 2013). Thrombin slowly accumulates during the amplification phase, activating platelets and platelet derived factor V, amplifying the prothrombinase activity. In a positive feedback loop, thrombin activates factor XI and VIII, generating more factor Xa. Hemophilia A and B are defined by deficiencies in factors VIII and IX, respectively, which is associated with spontaneous and prolonged bleeding. The extrinsic pathway can be assayed *in vitro* by measuring the prothrombin time (PT). The intrinsic pathway of coagulation can be triggered independently by collagen, polyphosphates secreted by platelets, neutrophil extracellular



**FIGURE 5**

Mitochondrial proteins upregulated during winter clustered using STRING. OXCT1 (Succinyl-CoA-3-ketoacid coenzyme A transferase 1), ACADL (Long-chain specific acyl-CoA dehydrogenase), PMPCB (Mitochondrial-processing peptidase subunit beta), MTX2 (Metaxin-2); GRPEL1 (GrpE protein homolog 1), UQCRC2 (Cytochrome b-c1 complex subunit 2), NDUFB10 (NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10), AMT (Aminomethyltransferase), ACAA2 (3-ketoacyl-CoA thiolase), PTC1 (Pentatricopeptide repeat-containing protein 1), IDH3 (Isocitrate dehydrogenase [NAD] subunit alpha), ECI1 (Enoyl-CoA delta isomerase 1), TSFM (Elongation factor Ts), ME2 (NAD-dependent malic enzyme), HCCS (Cytochrome c-type heme lyase), MRPL12 (39S ribosomal protein L12), NDUF2 (NADH dehydrogenase [ubiquinone] iron-sulfur protein 2), NUDT2 (Bis(5'-nucleosyl)-tetrphosphatase), HADHA (Trifunctional enzyme subunit alpha), IDH3B (Isocitrate dehydrogenase [NAD] subunit beta), PITRM1 (Presequence protease), AK3 (GTP-AMP phosphotransferase AK3), UQCRB (Cytochrome b-c1 complex subunit 7), NDUFAB1 (Acyl carrier protein).


traps (NETs), and artificial material such as glass, leading to activation of factors XII, XI, kallikrein and the subsequent downstream coagulation factors (Muller et al., 2009; Versteeg et al., 2013). The intrinsic pathway can be assessed *in vitro* by measuring the activated partial thromboplastin time (APTT).

## Secondary hemostasis adaptations during torpor

Secondary hemostasis is reduced during torpor in ground squirrels, bears, hedgehogs, and hamsters. In general, hibernating animals in torpor reduce the level of coagulation factors VIII, IX and XI resembling mild hemophilia A, B and C respectively (Denyes and Carter, 1961; Biorck et al., 1962; Lechler and Penick, 1963; Cooper et al., 2016; Welinder et al., 2016). The decreases in factor VIII, IX and XI during torpor in examined hibernating species are on average 78%, 61% and 51%, respectively, compared to euthermic non-hibernating values (Figure 7). Interestingly, the extrinsic pathway of the coagulation cascade remains fairly unaltered during

hibernation. A logical explanation could be that thrombosis due to stasis of blood should be prevented, but blood clotting due to tissue damage by an intruder or predator (activating the extrinsic pathway) may still be vital. An alternative explanation is that during torpor, damage to blood vessels is unlikely, so suppressing the extrinsic pathway is not necessary and protection against stasis-induced clots is more important. Remarkably, one clotting factor consistently showed an increase in protein levels during torpor: prothrombin. Possibly, this is caused by a decrease in its normal baseline formation resulting from inhibition of spontaneous, low-level activation of the clotting cascade, thus diminishing the conversion of prothrombin into thrombin (Lechler and Penick, 1963; Welinder et al., 2016). Prothrombin mRNA levels were slightly decreased in the livers of hibernating 13-lined ground squirrels, consistent with this explanation (Gillen et al., 2021). To assess the overall level of activation of clotting, levels of the irreversible complex between thrombin and its inhibitor antithrombin may be used. Thrombin-antithrombin complexes were decreased during hibernation in 13-lined ground squirrels, consistent with suppression of secondary hemostasis (Bonis et al., 2019). An alternative to reducing clotting






factors would be to increase anticoagulants. Yet, the anticoagulant proteins antithrombin and protein C are not reduced in torpid hamsters and ground squirrels (Bonis et al., 2019). However, in hibernating bears antithrombin levels are reduced, but it is not known if this is due to decreased production of antithrombin or increased consumption (Welinder et al., 2016). Collectively, torpor features the suppression of both the intrinsic and extrinsic arms of the coagulation cascade leading to prolonged APTT times and a mild extension of PT times (Biorck et al., 1962; Lechler and Penick, 1963; Pivorun and Sinnamon, 1981; De Wit et al., 1985; Iles et al., 2017). Diverse hibernators including shallow hibernators (bears), facultative hibernators (hamsters), and deep seasonal hibernators (ground squirrels and bats) use similar adaptations. These common strategies reduce both primary and secondary hemostasis leading to prolonged APTT (Figure 7).

The decreased activities of clotting factors in the blood correlate well with the changes in APTT and PT, however, the differences in protein levels could be due to decreased synthesis or increased consumption during torpor. Many of the blood proteins involved in secondary hemostasis are produced in liver, and a few are released from platelets and endothelial cells. In addition to traditional activity assays, transcriptome, proteome, and metabolomes are being analyzed in liver and plasma to identify changes in hemostatic regulators. The livers of torpid 13-lined ground squirrels revealed

decreases in prothrombin, factor V, factor IX, tissue factor, and heparin cofactor II mRNA while VWF mRNA was decreased in bone marrow (Cooper et al., 2016; Gillen et al., 2021). There was also an increase in  $\alpha 2$  macroglobulin mRNA in liver during hibernation, a serine protease inhibitor that can inhibit a broad range of clotting factors and its protein level has been previously shown to increase during torpor (Srere et al., 1995), potentially contributing to the suppressed state of hemostasis in torpor. Although these findings are in line with a suppression of hemostasis in torpor, some other studies demonstrate changes that are less unidirectional or involve both procoagulant and anticoagulant factors. For example, clotting factor mRNA expression was unexpectedly decreased in the brains of hibernating Himalayan marmots, but not in their livers (Bai et al., 2019). Another marmot transcriptome study compared two species, but did not look into seasonal variation in expression (Liu et al., 2016). While transcriptomes are useful in measuring new synthesis of proteins, activity in plasma is ultimately estimated by measuring levels of circulating protein which can be detected using proteomics. The bear plasma proteome revealed decreases in clotting factors similar to that reported for individual factor assays with winter increases in  $\alpha 2$  macroglobulin, potentially inhibiting thrombin, whereas thrombin itself and fibrinogen were also increased. In contrast, factors VIII and IX and VWF decreased in winter, in line with a suppression of hemostasis, but also the anticoagulant



factors antithrombin and protein C decreased (Welinder et al., 2016). An arctic ground squirrel liver proteome also revealed increased  $\alpha 2$  macroglobulin but a decrease in antithrombin during torpor (Shao et al., 2010). Finally, a liver proteome of 13-lined ground squirrels showed no seasonal differential expression of proteins involved in hemostasis (Rose et al., 2011). Although all analyzed hibernating mammals demonstrated some suppression within the hemostatic system, there remains some variation between species regarding which factors are involved and whether some procoagulant factors are increased. For the most part proteomic studies seem to align better with assays of individual proteins than do transcriptomic studies. This suggests that protein levels may be regulated more by consumption than transcriptionally. Further analysis of the plasma and liver proteomes may reveal other adaptations that regulate hemostasis.

## Fibrinolysis adaptations during torpor

Proteins released from endothelial cells after clot formation initiate fibrinolysis. The cross-linked fibrin network is

enzymatically degraded by plasmin, which is formed from plasminogen by tissue plasminogen activator (t-PA). t-PA is slowly released by damaged endothelium enabling a gradual degradation of fibrin after the bleeding has stopped and tissue regeneration has started (Gonias, 2021). A marker of fibrinolysis activation is the level of circulating complexes between tPA and its inhibitor plasminogen activator inhibitor 1 (PAI-1). In ground squirrel torpor, PAI-1 levels are decreased along with tPA-PAI-1 complexes consistent with a hyperfibrinolytic state (Bonis et al., 2019). In ground squirrels, plasminogen levels drop two-fold while in bears and hamsters an increase is seen during torpor. Upon fibrinolysis, fibrin in a clot is cleaved into fibrin degradation products, of which D-dimers can be detected in plasma and is commonly used in the diagnosis of venous or arterial thrombosis. In bears, hamsters, and ground squirrels no increases in D-dimers are seen during hibernation, consistent with low levels of clot formation and fibrinolysis (Iles et al., 2017; Bonis et al., 2019). The lack of D-dimers is consistent with suppression of secondary hemostasis and absence of actual clot fibrinolysis during torpor, while the hyperfibrinolytic state reflected by increased tPA and PAI-1 may be an added layer of protection in the event that a clot does form (Figure 7).

## Medical applications

The pathways that prevent activation and clearance of circulating platelets throughout hibernation, key receptors and ligands for platelet margination, and suppression of activation and apoptosis in the cold remain to be determined. Yet, unraveling these pathways could have direct applications in the storage of human platelets. Mimicking the hibernator's platelet resilience to extended cold storage by unlocking their cell-autonomous adaptations might allow long-term cold storage for human platelet transfusion, thus improving transfusion availability and reducing costs. Human red blood cells can be stored in the cold, and research on arctic ground squirrel red blood cells stored in the cold revealed novel protective mechanisms that may be applicable to human red blood cells, but is beyond the scope of this review on hemostasis and hibernation (Gehrke et al., 2019). Additionally, expanding our knowledge about the effects of lower temperature on hemostasis may help understand its role in patients with accidental hypothermia, e.g., near drowning, or those undergoing therapeutic hypothermia. Advancement of these insights will help physicians to better evaluate advantages and disadvantages of therapeutic hypothermia. More applications for therapeutic hypothermia may arise, as mild intraoperative hypothermia may for instance be beneficial in cardiac surgery (Hendriks et al., 2022) and plastic surgery by reducing thrombosis in free tissue transfer, hence improving free flap survival (Liu et al., 2011).

Understanding the mechanisms that hibernators use to prevent activation and clearance of platelets, and prevent thromboembolic complications, may be relevant for other medical applications other than surgery. Pulmonary embolism arising from DVT is a serious medical condition caused by immobilization leading to activation of blood clotting. A study in bears revealed the role of decreased platelet HSP47 in preventing clot formation (Thienel et al., 2023), and this could be translated to humans and other

TABLE 2 Secondary hemostasis in hibernation.

Measurement	Euthermia (EU)	Torpor	Arousal	Species	References
Thrombin time (sec)	13.2 ± 0.7	15.8 ± 0.7*	14.7 ± 1.3	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
	37°C	6°C	37°C		
PT (sec)	9.8 ± 1.6	10.6 ± 1.5		13-lined ground squirrel	Lechler and Penick (1963)
	36.3°C	7.9°C			
	8.1 ± 0.3	8.3 ± 0.2	14.0 ± 2.1 * <sup>#</sup>	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
	37°C	6°C	37°C		
	42.6	60.5	51.2	European Hedgehog	De Wit et al. (1985)
	8.75 ± 0.5			Golden hamster	Deveci et al. (2001)
	41–48	38–62		Red-eared slider (turtle); Painted turtle	Barone and Jacques (1975)
	8.6 ± 0.3	7.6 ± 1.5	8.1 ± 2.2	American black bear	Iles et al. (2017)
	22	23	22	Hedgehog	Biorck et al. (1962)
(%EU)	100% ± 3% (n = 5) 37°C	129% ± 16% * (n = 5) 20°C	110% ± 4% <sup>#</sup> (n = 5) 35°C	C57Bl/6 pharmacological torpor	(De Vrij et al. unpublished)
(%EU)	100% ± 9% (n = 10)	184% ± 82% (n = 3)	84% ± 6% <sup>#</sup> (n = 4)	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
APTT (sec)	45.5 ± 8.7	109.3 ± 42*		13-lined ground squirrel	Lechler and Penick (1963)
	36°C	8°C			
	25.0 ± 1.0	51.0 ± 2.4*	30.1 ± 1.4 <sup>#</sup>	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
	36°C	6°C	37°C		
	29.8 ± 0.27			Golden hamster	Deveci et al. (2001)
	17.6 ± 0.8	29.7 ± 7.9*	24.5 ± 2.1*	American black bear	Iles et al. (2017)
(%EU)	100% ± 22% (n = 10)	339% ± 88% (n = 3)	86% ± 12% <sup>#</sup> (n = 4)	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Thrombin generation (%EU)	100% ± 23% (n = 3) 36°C	8% ± 19%* (n = 7) 9°C	61% ± 58% (n = 5) 37°C	Syrian hamster	(De Vrij et al. unpublished)
Prothrombin (U/mL)	443 ± 132 36.3°C	698 ± 143* 7.9°C		13-lined ground squirrel	Lechler and Penick (1963)
(IU/mL)	1.10 ± 0.19	1.29 ± 0.31*		Scandinavian brown bear	Welinder et al. (2016)
(Sec)	11.5 ± 0.3 37°C	11.9 ± 0.3 6°C	10.5 ± 0.3* <sup>#</sup> 37°C	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
(U/mL)	20	55	90	Hedgehog	Biorck et al. (1962)
(%EU)	100% ± 10% 36°C	116% ± 20% 9°C	184% ± 15% * 37°C	Syrian hamster	(De Vrij et al. unpublished)
Residual prothrombin in serum (U/mL)	27 ± 38 36°C	449 ± 157* 8°C		13-lined ground squirrel	Lechler and Penick (1963)
Factor II, VII, X combined assay (sec)	12.8 ± 0.4 37°C	14.3 ± 0.6 6°C		Franklin's ground squirrel	Pivorun and Sinnamon (1981)
Factor V	639% ± 212% 36°C	570% ± 143%		13-lined ground squirrel	Lechler and Penick (1963)
		7.9°C			
(Sec)	16.2 ± 0.3	20.1 ± 0.3*	17.8 ± 0.4* <sup>#</sup> 37°C	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
	37°C	6°C			

(Continued on following page)

TABLE 2 (Continued) Secondary hemostasis in hibernation.

Measurement	Euthermia (EU)	Torpor	Arousal	Species	References
(% human plasma)	450	542	354	Hedgehog	Biorck et al. (1962)
(%EU)	100% ± 44%	20% ± 5% *	72% ± 26%	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Factor VII	369% ± 138% 36°C	536%		13-lined ground squirrel	Lechler and Penick (1963)
		8°C			
	1.01 ± 0.6	0.57 ± 0.14		Scandinavian brown bear	Welinder et al. (2016)
(%EU)	100% ± 15%	93% ± 22%	133% ± 11%	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Factor VIII	165% ± 73% 36°C	35% ± 11%*		13-lined ground squirrel	Lechler and Penick (1963)
		8°C			
(% relative to human plasma)	232 ± 2.0	68 ± 0.1*	230%	13-lined ground squirrel	Cooper et al. (2016)
(IU/mL)	2.92 ± 1.03	0.86 ± 0.35 *		Scandinavian brown bear	Welinder et al. (2016)
(%EU)	100% ± 15%	11% ± 5% *	52% ± 37%	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Factor IX (% human plasma)	378 ± 157	188 ± 65*		13-lined ground squirrel	Lechler and Penick (1963)
	36°C	8°C			
	425 ± 20	140 ± 4.0*	380%	13-lined ground squirrel	Cooper et al. (2016)
(%EU)	100% ± 13%	34% ± 10%	150% ± 23%*	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Factor X	867% ± 126% 36°C	805% ± 269% 8°C		13-lined ground squirrel	Lechler and Penick (1963)
(Sec)	18.9 ± 0.4	18.7 ± 0.7	19.4 ± 0.3	Franklin's ground squirrel	Pivorun and Sinnamon (1981)
	37°C	6°C	37°C		
(%EU)	100% ± 16%	131% ± 30%	138% ± 13%	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Factor XI (%)	111	72		13-lined ground squirrel	Lechler and Penick (1963)
	36°C	8°C			
(%EU)	100% ± 22%	34% ± 14% *	92% ± 31% <sup>‡</sup>	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Factor XII	291% ± 71% 36°C	222% ± 70%		13-lined ground squirrel	Lechler and Penick (1963)
		8°C			
Fibrinogen (mg%)	189 ± 49	145 ± 34		13-lined ground squirrel	Lechler and Penick (1963)
	36°C	8°C			
(g/L)	2.09 ± 0.94	2.26 ± 0.46		Scandinavian brown bear	Welinder et al. (2016)
(%)	0.54	0.5	0.29	Hedgehog	Biorck et al. (1962)
(%EU)	100% ± 20%	45% ± 20%	130% ± 65% <sup>‡</sup>	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Plasminogen (%EU)	100% ± 31% (n = 10)	168% ± 51% (n = 5)	249% ± 51% (n = 5) *		(De Vrij et al. unpublished)

(Continued on following page)

TABLE 2 (Continued) Secondary hemostasis in hibernation.

Measurement	Euthermia (EU)	Torpor	Arousal	Species	References
(% relative to human plasma)	37.6% ± 9.2%	19.7% ± 12.3%		13-lined ground squirrel	Bonis et al. (2019)
PAI-1 (ng/mL)	2.15 ± 1.04	0.45 ± 0.35		13-lined ground squirrel	Bonis et al. (2019)
Plasmin inhibitor (%EU)	100% ± 5%	89% ± 18%	109% ± 24%	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Protein C (IU/mL)	0.44 ± 0.08	0.33 ± 0.08*		Scandinavian brown bear	Welinder et al. (2016)
(%EU)	100% ± 12%	61% ± 9%	112% ± 14%*	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Antithrombin (IU/mL)	0.98 ± 0.09	0.47 ± 0.04*		Scandinavian brown bear	Welinder et al. (2016)
(%EU)	100% ± 6%	86% ± 11%	104% ± 7.2%	Syrian hamster	(De Vrij et al. unpublished)
	36°C	9°C	37°C		
Heparin (sec)	36.6 ± 0.3 37°C	37.7 ± 0.4 6°C		Franklin's ground squirrel	Pivorun and Sinnamon (1981)
D-dimer (ug/L)		152	124	American black bear	Iles et al. (2017)
	15 ± 13	28 ± 32	10 ± 17	Syrian hamster	de Vrij et al. (2021)
	36.3°C ± 0.9°C	8.8°C ± 0.7°C	36.7°C ± 1.1°C		
(%EU)	100% ± 55%	76% ± 22%	80% ± 26%	13-lined ground squirrel	Bonis et al. (2019)


mammals under immobilization. This research could have direct applications in screening patients at risk of DVT with elevated HSP47 and also by developing therapeutic treatments to block HSP47 in immobilized patients. Diffuse intravascular coagulation (DIC) is a severe complication that can arise in critically ill patients. Sepsis, a syndrome of organ failure due to a dysregulated host response to infection, is a common cause of critical illness and DIC. DIC is associated with thromboembolic complications on the one hand and bleeding complications due to consumption of platelets and coagulation factors on the other hand. Therapeutic hypothermia may improve the coagulopathy as measured by thromboelastography in patients with sepsis or septic shock (Johansen et al., 2015), and even decrease mortality and end-organ damage in sepsis as demonstrated experimentally (Acosta-Lara and Varon, 2013). Preventing fever and maintaining normothermia (37°C) by cooling volunteers upon injection of endotoxin, decreases markers of DIC that were induced by endotoxin challenge (Harmon et al., 2021). Yet, therapeutic hypothermia did not lower mortality in sepsis (Itenov et al., 2018).

The use of anticoagulant drugs to prevent thromboembolic complications in high-risk situations after major surgery, in patients with atrial fibrillation, or after a previous unproven thromboembolic event, is associated with bleeding complications as major side effect. Moreover, bleeding is the most common adverse drug event bringing patients to emergency wards (Leendertse et al., 2008; Budnitz et al., 2011) and the newest class of anticoagulants, direct oral anticoagulants (DOAC), are associated with fatal bleeding on 0.16 per 100 patient-years (Chai-Adisaksopha et al., 2015). Therefore bleeding requires correction of anticoagulation as quickly as possible. If a pharmacological tool becomes available to

mimic the torpor induced suppression of thrombosis, it may add the benefit of a rapid reversal strategy, since arousal rapidly reverses antithrombotic effects within minutes to hours. However, most reversal techniques require several hours to reverse anticoagulation, i.e., by administering fresh frozen plasma, Vitamin K or prothrombin complex concentrate (Fredriksson et al., 1992; Hanley, 2004). To date, only two registered monoclonal antibody fragments against DOAC are faster and reverse anticoagulation within minutes (White et al., 2022), but are highly expensive. Therefore, the development of new treatment and reversal strategies may help physicians to effectively manage the life-threatening emergency of bleeding in anticoagulated patients (Christos and Naples, 2016). Unraveling the mechanisms driving the hemostatic adaptations in hibernators might contribute to the development of these novel, reversible and safer anticoagulants with lower risk on coagulation abnormalities—even at euthermia—in patients with critical illnesses like sepsis, trauma and after major surgery.

## Summary and future research

In spite of the expected risk of thrombosis during hibernation due to prolonged immobility with reduced blood flow and increased blood viscosity, hibernating mammals do not show signs of thrombotic complications during or after hibernation (Figure 8). During torpor in diverse hibernating species of mammals, there is a rapid and reversible anti-thrombotic shift by the reduction in circulating platelets, VWF, and coagulation factors. This results in a mixed phenotype resembling



thrombocytopenia, Von Willebrand Disease, and Hemophilia A, B and C. At the same time, torpid animals maintain factors required for fibrinolysis. One of the most remarkable adaptations is the ability to store platelets in the liver sinusoids during torpor, releasing them intact after prolonged periods of storage in the cold. Future research should use phosphoproteome and metabolomics studies to identify seasonal differences in signaling pathways and molecules released by platelets, in addition to characterizing mitochondrial function. Many common adaptations protect hibernating mammals from unwanted activation of hemostasis during torpor and yet allow for a quick reversal of this state to restore proper hemostasis during arousal. The mechanisms behind these reversible adaptations remain to be discovered and may open doors for novel therapies and improved treatments.

## Author contributions

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

## Funding

S.T.C. received funding from the National Institutes of Health, USA.

## Acknowledgments

We would like to thank T. Lisman and J. Adelmeijer from the Surgical Research Laboratory, Department of Surgery, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands, as well as M. Lukens and R. Mulder from the Department of Laboratory Medicine, University Medical Center Groningen, Groningen, the Netherlands, for their help in measurements from hibernating Syrian hamster and interpretation of these data.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Acosta-Lara, P., and Varon, J. (2013). Therapeutic hypothermia in sepsis: To use or not to use? *Am. J. Emerg. Med.* 31, 381–382. doi:10.1016/j.ajem.2012.09.017
- Ahmad, N., Dube, B., Agarwal, G. P., and Dube, R. K. (1979). Comparative studies of blood coagulation in hibernating and non-hibernating frogs (*Rana tigrina*). *Thromb. Haemost.* 42, 959–964. doi:10.1055/s-0038-1656986
- Arinell, K., Blanc, S., Welinder, K. G., Støen, O. G., Evans, A. L., and Frøbert, O. (2018). Physical inactivity and platelet function in humans and Brown bears: A comparative study. *Platelets* 29, 87–90. doi:10.1080/09537104.2017.1336530
- Bai, L., Liu, B., Ji, C., Zhao, S., Liu, S., Wang, R., et al. (2019). Hypoxic and cold adaptation insights from the himalayan marmot genome. *iScience* 11, 519–530. doi:10.1016/j.isci.2018.11.034
- Barone, M. C., and Jacques, F. A. (1975). The effects of induced cold torpor and time of year on blood coagulation in *Pseudemys scripta elegans* and *Chrysemys picta bellii*. *Comp. Biochem. Physiol. A Comp. Physiol.* 50, 717–721. doi:10.1016/0300-9629(75)90134-6
- Becker, G. A., Tuccelli, M., Kunicki, T., Chalos, M. K., and Aster, R. H. (1973). Studies of platelet concentrates stored at 22°C nad 4°C. *Transfusion* 13, 61–68. doi:10.1111/j.1537-2995.1973.tb05442.x
- Berger, G., Hartwell, D. W., and Wagner, D. D. (1998). P-Selectin and platelet clearance. *Blood* 92, 4446–4452. doi:10.1182/blood.v92.11.4446
- Björck, G., Johansson, B. W., and Nilsson, I. M. (1962). Blood coagulation studies in hedgehogs, in a hibernating and a non-hibernating state, and in dogs, hypothermic and normothermic. *Acta Physiol. Scand.* 56, 334–348. doi:10.1111/j.1748-1716.1962.tb02510.x
- Bonis, A., Anderson, L., Talhouarne, G., Schueller, E., Unke, J., Krus, C., et al. (2019). Cardiovascular resistance to thrombosis in 13-lined ground squirrels. *J. Comp. Physiol. B* 189, 167–177. doi:10.1007/s00360-018-1186-x
- Bouma, H. R., Strijkstra, A. M., Boerema, A. S., Deelman, L. E., Epema, A. H., Hut, R. A., et al. (2010). Blood cell dynamics during hibernation in the European Ground Squirrel. *Vet. Immunol. Immunopathol.* 136, 319–323. doi:10.1016/j.vetimm.2010.03.016
- Budnitz, D. S., Lovegrove, M. C., Shehab, N., and Richards, C. L. (2011). Emergency hospitalizations for adverse drug events in older Americans. *N. Engl. J. Med.* 365, 2002–2012. doi:10.1056/NEJMsa1103053
- Carey, H. V., Andrews, M. T., and Martin, S. L. (2003). Mammalian hibernation: Cellular and molecular responses to depressed metabolism and low temperature. *Physiol. Rev.* 83, 1153–1181. doi:10.1152/physrev.00008.2003
- Chai-Adisaksopha, C., Hillis, C., Isayama, T., Lim, W., Iorio, A., and Crowther, M. (2015). Mortality outcomes in patients receiving direct oral anticoagulants: A systematic review and meta-analysis of randomized controlled trials. *J. Thromb. Haemost.* 13, 2012–2020. doi:10.1111/jth.13139
- Chan, K. M., and Beard, K. (1993). A patient with recurrent hypothermia associated with thrombocytopenia. *Postgrad. Med. J.* 69, 227–229. doi:10.1136/pgmj.69.809.227
- Christos, S., and Naples, R. (2016). Anticoagulation reversal and treatment strategies in major bleeding: Update 2016. *West J. Emerg. Med.* 17, 264–270. doi:10.5811/westjem.2016.3.29294
- Cooper, S., Lloyd, S., Koch, A., Lin, X., Dobbs, K., Theisen, T., et al. (2017). Temperature effects on the activity, shape, and storage of platelets from 13-lined ground squirrels. *J. Comp. Physiol. B* 187, 815–825. doi:10.1007/s00360-017-1081-x
- Cooper, S., Sell, S., Nelson, L., Hawes, J., Benrud, J. A., Kohlhofer, B. M., et al. (2016). Von Willebrand factor is reversibly decreased during torpor in 13-lined ground squirrels. *J. Comp. Physiol. B* 186, 131–139. doi:10.1007/s00360-015-0941-5
- Cooper, S., Wilmarth, P. A., Cunliffe, J. M., Klimek, J., Pang, J., Tassi Yunga, S., et al. (2021). Platelet proteome dynamics in hibernating 13-lined ground squirrels. *Physiol. Genomics* 53, 473–485. doi:10.1152/physiolgenomics.00078.2021
- Cooper, S. T., Richters, K. E., Melin, T. E., Liu, Z. J., Hordyk, P. J., Benrud, R. R., et al. (2012). The hibernating 13-lined ground squirrel as a model organism for potential cold storage of platelets. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 302, R1202–R1208. doi:10.1152/ajpregu.00018.2012
- Cooper, S. T., Sell, S. S., Fahrenkrog, M., Wilkinson, K., Howard, D. R., Bergen, H., et al. (2016). Effects of hibernation on bone marrow transcriptome in thirteen-lined ground squirrels. *Physiol. Genomics* 48, 513–525. doi:10.1152/physiolgenomics.00120.2015
- D'Alessandro, A., Liunbruno, G., Grazzini, G., and Zolla, L. (2010). Red blood cell storage: The story so far. *Blood Transfus.* 8, 82–88. doi:10.2450/2009.0122-09
- de Vrij, E. L., Bouma, H. R., Goris, M., Weerman, U., de Groot, A. P., Kuipers, J., et al. (2021). Reversible thrombocytopenia during hibernation originates from storage and release of platelets in liver sinusoids. *J. Comp. Physiol. B* 191, 603–615. doi:10.1007/s00360-021-01351-3
- de Vrij, E. L., Vogelaar, P. C., Goris, M., Houwertjes, M. C., Herwig, A., Dugbartey, G. J., et al. (2014). Platelet dynamics during natural and pharmacologically induced torpor and forced hypothermia. *PLoS One* 9, e93218. doi:10.1371/journal.pone.0093218
- De Wit, C. A., Persson, G., Nilsson, I. M., and Johansson, B. W. (1985). Circannual changes in blood coagulation factors and the effect of warfarin on the hedgehog *Erinaceus europaeus*. *Comp. Biochem. Physiol. A* 80, 43–47. doi:10.1016/0300-9629(85)90675-9
- DeLong, J. P., Gibert, J. P., Lühring, T. M., Bachman, G., Reed, B., Neyer, A., et al. (2017). The combined effects of reactant kinetics and enzyme stability explain the temperature dependence of metabolic rates. *Ecol. Evol.* 7, 3940–3950. doi:10.1002/ece3.2955
- Denyes, A., and Carter, J. D. (1961). Clotting-time of cold-exposed and hibernating hamsters. *Nature* 190, 450–451. doi:10.1038/190450a0
- Deveci, D., Stone, P. C., and Egginton, S. (2001). Differential effect of cold acclimation on blood composition in rats and hamsters. *J. Comp. Physiol. B* 171, 135–143. doi:10.1007/s003600000156
- Duguid, H., Simpson, R. G., and Stowers, J. M. (1961). Accidental hypothermia. *Lancet* 2, 1213–1219. doi:10.1016/s0140-6736(61)92588-0
- Egidi, M. G., D'Alessandro, A., Mandarello, G., and Zolla, L. (2010). Troubleshooting in platelet storage temperature and new perspectives through proteomics. *Blood Transfus.* 8 (3), s73–s81. doi:10.2450/2010.012S
- Fernandez-Moure, J. S., Van Eps, J. L., Cabrera, F. J., Barbosa, Z., Medrano Del Rosal, G., Weiner, B. K., et al. (2017). Platelet-rich plasma: A biomimetic approach to enhancement of surgical wound healing. *J. Surg. Res.* 207, 33–44. doi:10.1016/j.jss.2016.08.063
- Fredriksson, K., Norrving, B., and Strömblad, L. G. (1992). Emergency reversal of anticoagulation after intracerebral hemorrhage. *Stroke* 23, 972–977. doi:10.1161/01.str.23.7.972
- Friedrich, A. U., Kakuturu, J., Schnorr, P. J., Beyer, D. E., Jr., Palesty, J. A., Dickson, E. W., et al. (2017). Comparative coagulation studies in hibernating and summer-active black bears (*Ursus americanus*). *Thromb. Res.* 158, 16–18. doi:10.1016/j.thromres.2017.07.034
- Frøbert, O., Christensen, K., Fahlman, A., Brunberg, S., Josefsson, J., Särndahl, E., et al. (2010). Platelet function in Brown bear (*Ursus arctos*) compared to man. *Thromb. J.* 8, 11. doi:10.1186/1477-9560-8-11
- Gehrke, S., Rice, S., Stefanoni, D., Wilkerson, R. B., Nemkov, T., Reisz, J. A., et al. (2019). Red blood cell metabolic responses to torpor and arousal in the hibernator arctic ground squirrel. *J. Proteome Res.* 18, 1827–1841. doi:10.1021/acs.jproteome.9b00018
- Gillen, A. E., Fu, R., Riemondy, K. A., Jager, J., Fang, B., Lazar, M. A., et al. (2021). Liver transcriptome dynamics during hibernation are shaped by a shifting balance between transcription and RNA stability. *Front. Physiology* 12, 662132. doi:10.3389/fphys.2021.662132
- Gonias, S. L. (2021). Plasminogen activator receptor assemblies in cell signaling, innate immunity, and inflammation. *Am. J. Physiol. Cell Physiol.* 321, C721–c734. doi:10.1152/ajpcell.00269.2021
- Halikas, G., and Bowers, K. (1973). Seasonal variation in blood viscosity of the hibernating arctic ground squirrel (*Spermophilus undulatus plesius*). *Comp. Biochem. Physiol. A Comp. Physiol.* 44, 677–681. doi:10.1016/0300-9629(73)90522-7
- Hanley, J. P. (2004). Warfarin reversal. *J. Clin. Pathol.* 57, 1132–1139. doi:10.1136/jcp.2003.008904
- Harmon, M. B. A., Heijnen, N. F. L., de Bruin, S., Sperna Weiland, N. H., Meijers, J. C. M., de Boer, A. M., et al. (2021). Induced normothermia ameliorates the procoagulant host response in human endotoxaemia. *Br. J. Anaesth.* 126, 1111–1118. doi:10.1016/j.bja.2021.02.033
- Hendriks, K. D. W., Castela Forte, J. N., Kok, W. F., Mungroop, H. E., Bouma, H. R., Scheeren, T. W. L., et al. (2022). Mild hypothermia during cardiopulmonary bypass assisted CABG is associated with improved short- and long-term survival, a 18-year cohort study. *PLOS ONE* 17, e0273370. doi:10.1371/journal.pone.0273370
- Hendriks, K. D. W., Joschko, C. P., Hoogstra-Berends, F., Heegsma, J., Faber, K. N., and Henning, R. H. (2020). Hibernator-derived cells show superior protection and survival in hypothermia compared to non-hibernator cells. *Int. J. Mol. Sci.* 21, 1864. doi:10.3390/ijms21051864
- Hoffmeister, K. M., Felbinger, T. W., Falet, H., Denis, C. V., Bergmeier, W., Mayadas, T. N., et al. (2003). The clearance mechanism of chilled blood platelets. *Cell* 112, 87–97. doi:10.1016/s0092-8674(02)01253-9
- Hoffmeister, K. M. (2011). The role of lectins and glycans in platelet clearance. *J. Thromb. Haemost.* 9 (1), 35–43. doi:10.1111/j.1538-7836.2011.04276.x
- Horwitz, B. A., Chau, S. M., Hamilton, J. S., Song, C., Gorgone, J., Saenz, M., et al. (2013). Temporal relationships of blood pressure, heart rate, baroreflex function, and body temperature change over a hibernation bout in Syrian hamsters. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 305, R759–R768. doi:10.1152/ajpregu.00450.2012
- Hu, H. X., Du, F. Y., Fu, W. W., Jiang, S. F., Cao, J., Xu, S. H., et al. (2017). A dramatic blood plasticity in hibernating and 14-day hindlimb unloading Daurian ground squirrels (*Spermophilus dauricus*). *J. Comp. Physiol. B* 187, 869–879. doi:10.1007/s00360-017-1092-7
- Iles, T. L., Laske, T. G., Garshelis, D. L., and Iaizzo, P. A. (2017). Blood clotting behavior is innately modulated in *Ursus americanus* during early and late denning relative to summer months. *J. Exp. Biol.* 220, 455–459. doi:10.1242/jeb.141549
- Itenov, T. S., Johansen, M. E., Bestle, M., Thormar, K., Hein, L., Gyldensted, L., et al. (2018). Induced hypothermia in patients with septic shock and respiratory failure (CASS): A randomised, controlled, open-label trial. *Lancet Respir. Med.* 6, 183–192. doi:10.1016/S2213-2600(18)30004-3
- Jacobs, S. E., Berg, M., Hunt, R., Tarnow-Mordi, W. O., Inder, T. E., and Davis, P. G. (2013). Cooling for newborns with hypoxic ischaemic encephalopathy. *Cochrane Database Syst. Rev.* 2013, CD003311. doi:10.1002/14651858.CD003311

- Johansen, M. E., Jensen, J. U., Bestle, M. H., Ostrowski, S. R., Thormar, K., Christensen, H., et al. (2015). Mild induced hypothermia: Effects on sepsis-related coagulopathy—results from a randomized controlled trial. *Thromb. Res.* 135, 175–182. doi:10.1016/j.thromres.2014.10.028
- Kirkebo, A. (1968). Temperature effects on the viscosity of blood and the aorta distension from a hibernator, *Erinaceus europaeus* L. *Acta Physiol. Scand.* 73, 385–393. doi:10.1111/j.1365-201x.1968.tb10877.x
- Leblanc, R., and Peyruchaud, O. (2016). Metastasis: New functional implications of platelets and megakaryocytes. *Blood* 128, 24–31. doi:10.1182/blood-2016-01-636399
- Lebois, M., and Josefsson, E. C. (2016). Regulation of platelet lifespan by apoptosis. *Platelets* 27, 497–504. doi:10.3109/09537104.2016.1161739
- Lechler, E., and Penick, G. D. (1963). Blood clotting defect in hibernating ground squirrels (*Citellus tridecemlineatus*). *Am. J. Physiol.* 205, 985–988. doi:10.1152/ajplegacy.1963.205.5.985
- Leendertse, A. J., Egberts, A. C., Stoker, L. J., and van den Bemt, P. M.HARM Study Group (2008). Frequency of and risk factors for preventable medication-related hospital admissions in The Netherlands. *Arch. Intern. Med.* 168, 1890–1896. doi:10.1001/archinternmed.2008.3
- Liu, Y., Wang, B., Wang, L., Vikash, V., Wang, Q., Roggendorf, M., et al. (2016). Transcriptome analysis and comparison of *Marmota monax* and *Marmota himalayana*. *PLoS One* 11, e0165875. doi:10.1371/journal.pone.0165875
- Liu, Y. J., Hirsch, B. P., Shah, A. A., Reid, M. A., and Thomson, J. G. (2011). Mild intraoperative hypothermia reduces free tissue transfer thrombosis. *J. Reconstr. Microsurg* 27, 121–126. doi:10.1055/s-0030-1268211
- Lyman, C. P., and O'Brien, R. C. (1960). Circulatory changes in the thirteen-lined ground squirrel during the hibernation cycle. *Bull. Mus. Comp. Zoology* 124, 353–372.
- Mahajan, S. L., Myers, T. J., and Baldini, M. G. (1981). Disseminated intravascular coagulation during rewarming following hypothermia. *JAMA* 245, 2517–2518. doi:10.1001/jama.1981.03310490035022
- Mallet, M. L. (2002). Pathophysiology of accidental hypothermia. *QJM* 95, 775–785. doi:10.1093/qjmed/95.12.775
- Martin, S. L. (2008). Mammalian hibernation: A naturally reversible model for insulin resistance in man? *Diab Vasc. Dis. Res.* 5, 76–81. doi:10.3132/dvdr.2008.013
- McArthur, M. D., and Milsom, W. K. (1991). Changes in ventilation and respiratory sensitivity associated with hibernation in columbian (*Spermophilus columbianus*) and golden-mantled (*Spermophilus lateralis*) ground squirrels. *Physiol. Zool.* 64, 940–959. doi:10.1086/physzool.64.4.30157950
- Michelson, A. D., Barnard, M. R., Khuri, S. F., Rohrer, M. J., MacGregor, H., and Valeri, C. R. (1999). The effects of aspirin and hypothermia on platelet function *in vivo*. *Br. J. Haematol.* 104, 64–68. doi:10.1046/j.1365-2141.1999.01146.x
- Mikhailidis, D. P., and Barradas, M. A. (1993). A patient with recurrent hypothermia associated with thrombocytopenia. *Postgrad. Med. J.* 69, 752. doi:10.1136/pgmj.69.815.752
- Morrell, C. N., Murata, K., Swaim, A. M., Mason, E., Martin, T. V., Thompson, L. E., et al. (2008). *In vivo* platelet-endothelial cell interactions in response to major histocompatibility complex alloantibody. *Circ. Res.* 102, 777–785. doi:10.1161/CIRCRESAHA.107.170332
- Muller, F., Mutch, N. J., Schenk, W. A., Smith, S. A., Esterl, L., Spronk, H. M., et al. (2009). Platelet polyphosphates are proinflammatory and procoagulant mediators *in vivo*. *Cell* 139, 1143–1156. doi:10.1016/j.cell.2009.11.001
- Nieswandt, B., Pleines, I., and Bender, M. (2011). Platelet adhesion and activation mechanisms in arterial thrombosis and ischaemic stroke. *J. Thromb. Haemost.* 9 (1), 92–104. doi:10.1111/j.1538-7836.2011.04361.x
- Ou, J., Ball, J. M., Luan, Y., Zhao, T., Miyagishima, K. J., Xu, Y., et al. (2018). iPSCs from a hibernator provide a platform for studying cold adaptation and its potential medical applications. *Cell* 173, 851–863. doi:10.1016/j.cell.2018.03.010
- Öztop, M., Özbek, M., Liman, N., Beyaz, F., Ergün, E., and Ergün, L. (2019). Localization profiles of natriuretic peptides in hearts of pre-hibernating and hibernating Anatolian ground squirrels (*Spermophilus xanthoprimum*). *Vet. Res. Commun.* 43, 45–65. doi:10.1007/s11259-019-9745-5
- Peerschke, E. I., Yin, W., and Ghebrehiwet, B. (2008). Platelet mediated complement activation. *Adv. Exp. Med. Biol.* 632, 81–91. doi:10.1007/978-0-387-78952-1\_7
- Pivorun, E. B., and Sinnamon, W. B. (1981). Blood coagulation studies in normothermic, hibernating, and aroused *Spermophilus franklini*. *Cryobiology* 18, 515–520. doi:10.1016/0011-2240(81)90212-1
- Polderman, K. H. (2009). Mechanisms of action, physiological effects, and complications of hypothermia. *Crit. Care Med.* 37, S186–S202. doi:10.1097/CCM.0b013e3181aa5241
- Rashid, N., Irfan, M., Nadeem, M. S., and Shabbir, A. (2016). Comparative seasonal haematology of two bat species, *scotophilus heathii* and *Pipistrellus pipistrellus*, in a subtropical area of Pakistan. *Pak. J. Zoology* 48, 1503.
- Reddick, R. L., Poole, B. L., and Penick, G. D. (1973). Thrombocytopenia of hibernation. Mechanism of induction and recovery. *Lab. Invest.* 28, 270–278.
- Rohrer, M. J., and Natale, A. M. (1992). Effect of hypothermia on the coagulation cascade. *Crit. Care Med.* 20, 1402–1405. doi:10.1097/00003246-199210000-00007
- Rose, J. C., Epperson, L. E., Carey, H. V., and Martin, S. L. (2011). Seasonal liver protein differences in a hibernator revealed by quantitative proteomics using whole animal isotopic labeling. *Comp. Biochem. Physiol. Part D. Genomics Proteomics* 6, 163–170. doi:10.1016/j.cbd.2011.02.003
- Semple, J. W., Italiano, J. E., Jr., and Freedman, J. (2011). Platelets and the immune continuum. *Nat. Rev. Immunol.* 11, 264–274. doi:10.1038/nri2956
- Shao, C., Liu, Y., Ruan, H., Li, Y., Wang, H., Kohl, F., et al. (2010). Shotgun proteomics analysis of hibernating arctic ground squirrels. *Mol. Cell. Proteomics MCP* 9, 313–326. doi:10.1074/mcp.M900260-MCP200
- Splinter, N., Mancosky, A., Laffin, C., Clement, M., Nisius, M., Arbs, B., et al. (2023). Platelets from 13lined ground squirrels are resistant to cold storage lesions. *J. Comp. Physiol. B.* 193 (1), 125–134. doi:10.1007/s00360-022-01469-y
- Srere, H. K., Belke, D., Wang, L. C., and Martin, S. L. (1995). Alpha 2-Macroglobulin gene expression during hibernation in ground squirrels is independent of acute phase response. *Am. J. Physiol.* 268, R1507–R1512. doi:10.1152/ajpregu.1995.268.6.R1507
- Stine, R. J. (1977). Accidental hypothermia. *JACEP* 6, 413–416. doi:10.1016/s0361-1124(77)80007-5
- Stolla, M., Bailey, S. L., Fang, L., Fitzpatrick, L., Gettinger, I., Pellham, E., et al. (2020). Effects of storage time prolongation on *in vivo* and *in vitro* characteristics of 4°C-stored platelets. *Transfusion* 60, 613–621. doi:10.1111/trf.15669
- Svihla, A., Bowman, H. R., and Ritenour, R. (1951). Prolongation of clotting time in dormant estivating mammals. *Science* 114, 298–299. doi:10.1126/science.114.2960.298
- Talaei, F., Bouma, H. R., Hylkema, M. N., Strijkstra, A. M., Boerema, A. S., Schmidt, M., et al. (2012). The role of endogenous H2S formation in reversible remodeling of lung tissue during hibernation in the Syrian hamster. *J. Exp. Biol.* 215, 2912–2919. doi:10.1242/jeb.067363
- Talaei, F., Bouma, H. R., Van der Graaf, A. C., Strijkstra, A. M., Schmidt, M., and Henning, R. H. (2011). Serotonin and dopamine protect from hypothermia/rewarming damage through the CBS/H2S pathway. *PLoS One* 6, e22568. doi:10.1371/journal.pone.0022568
- Thienel, M., Müller-Reif, J. B., Zhang, Z., Ehreiser, V., Huth, J., Shchurovska, K., et al. (2023). Immobility-associated thromboprotection is conserved across mammalian species from bear to human. *Science* 380, 178–187. doi:10.1126/science.abo5044
- Utz, J. C., Nelson, S., O'Toole, B. J., and van Breukelen, F. (2009). Bone strength is maintained after 8 months of inactivity in hibernating golden-mantled ground squirrels, *Spermophilus lateralis*. *J. Exp. Biol.* 212, 2746–2752. doi:10.1242/jeb.032854
- Valeri, C. R., MacGregor, H., Cassidy, G., Tinney, R., and Pompei, F. (1995). Effects of temperature on bleeding time and clotting time in normal male and female volunteers. *Crit. Care Med.* 23, 698–704. doi:10.1097/00003246-199504000-00019
- Van Der Wal, D. E., Du, V. X., Lo, K. S. L., Rasmussen, J. T., Verhoef, S., and Akkerman, J. W. N. (2010). Platelet apoptosis by cold-induced glycoprotein Iba clustering. *J. Thrombosis Haemostasis* 8, 2554–2562. doi:10.1111/j.1538-7836.2010.04043.x
- Van Poucke, S., Stevens, K., Marcus, A. E., and Lance, M. (2014). Hypothermia: Effects on platelet function and hemostasis. *Thromb. J.* 12, 31. doi:10.1186/s12959-014-0031-z
- Vella, M. A., Jenner, C., Betteridge, D. J., and Jowett, N. I. (1988). Hypothermia-induced thrombocytopenia. *J. R. Soc. Med.* 81, 228–229. doi:10.1177/014107688808100414
- Versteeg, H. H., Heemsker, J. W., Levi, M., and Reitsma, P. H. (2013). New fundamentals in hemostasis. *Physiol. Rev.* 93, 327–358. doi:10.1152/physrev.00016.2011
- Wang, C. H., Chen, N. C., Tsai, M. S., Yu, P. H., Wang, A. Y., Chang, W. T., et al. (2015). Therapeutic hypothermia and the risk of hemorrhage: A systematic review and meta-analysis of randomized controlled trials. *Med. Baltim.* 94, e2152. doi:10.1097/MD.0000000000002152
- Ware, J., Corken, A., and Khetpal, R. (2013). Platelet function beyond hemostasis and thrombosis. *Curr. Opin. Hematol.* 20, 451–456. doi:10.1097/MOH.0b013e32836344d3
- Welinder, K. G., Hansen, R., Overgaard, M. T., Brohus, M., Sønderkær, M., von Bergen, M., et al. (2016). Biochemical foundations of health and energy conservation in hibernating free-ranging subadult Brown bear *Ursus arctos*. *J. Biol. Chem.* 291, 22509–22523. doi:10.1074/jbc.M116.742916
- White, J. G., and Krivit, W. (1967). An ultrastructural basis for the shape changes induced in platelets by chilling. *Blood* 30, 625–635. doi:10.1182/blood.v30.5.625.625
- White, K., Faruqi, U., and Cohen, A. A. T. (2022). New agents for DOAC reversal: A practical management review. *Br. J. Cardiol.* 29, 1. doi:10.5837/bjc.2022.001
- Winokur, R., and Hartwig, J. H. (1995). Mechanism of shape change in chilled human platelets. *Blood* 85, 1796–1804. doi:10.1182/blood.v85.7.1796.bloodjournal8571796
- Xavier, R. G., White, A. E., Fox, S. C., Wilcox, R. G., and Heptinstall, S. (2007). Enhanced platelet aggregation and activation under conditions of hypothermia. *Thromb. Haemost.* 98, 1266–1275. doi:10.1160/th07-03-0189
- Zatzman, M. L. (1984). Renal and cardiovascular effects of hibernation and hypothermia. *Cryobiology* 21, 593–614. doi:10.1016/0011-2240(84)90220-7